

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
17 October 2002 (17.10.2002)

PCT

(10) International Publication Number
WO 02/080913 A1

- (51) International Patent Classification⁷: **A61K 31/425**,
A61P 35/00, C07D 417/12, 417/14
- (21) International Application Number: PCT/EP02/03746
- (22) International Filing Date: 4 April 2002 (04.04.2002)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
01107820.1 6 April 2001 (06.04.2001) EP
- (71) Applicant (*for all designated States except US*): **F. HOFFMANN-LA ROCHE AG** [CH/CH]; Grenzacherstrasse 124, CH-4070 Basle (CH).
- (72) Inventors; and
- (75) Inventors/Applicants (*for US only*): **BAUER, Sabine** [DE/DE]; Primelstrasse 3a, 82377 Penzberg (DE). **MEYER, Markus** [DE/DE]; Maurenweg 6, 79395 Neuenburg (DE). **SCHNITZER, Tobias** [DE/DE]; Bellizonastrasse 5, 81475 Muenchen (DE). **SCHUELL, Christine** [DE/DE]; Wiesenpromenade West 42, 64673 Zwingenberg (DE). **WOLFF, Hans-Peter** [DE/DE]; Huegelstrasse 11, 69469 Weinheim (DE).
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:**
— *with international search report*
— *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments*
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: THIAZOLIDINEDIONES ALONE OR IN COMBINATION WITH OTHER THERAPEUTIC AGENTS FOR INHIBITING OR REDUCING TUMOUR GROWTH

(57) Abstract: Use of thiazolidinedione derivatives for the preparation of medicaments for inhibiting or reducing tumour growth or metastases, alone or in combination with an RXR agonist or well-known antitumour agent.

WO 02/080913 A1

**Thiazolidinediones alone or in combination with other therapeutic agents
for inhibiting or reducing tumour growth**

Field of the invention

5 The present invention relates to ligands that bind to and affect the peroxisome proliferator-activated receptor (PPAR) gamma as well as their therapeutic use alone or in combination with other therapeutic agents for inhibiting or reducing the growth of tumours.

Background of the invention

10 The peroxisome proliferator-activated receptors (PPARs) belong to the steroid receptor superfamily and, as such, are ligand activated transcription factors and exist in different subtypes and isoforms (see, for example, Pershadsingh, H.A., Exp. Opin. Invest. Drugs 8 (1999) 1859-1872; Willson, T.M., et al., J. Med. Chem. 43 (2000) 527-550; Kersten, S., et al., Nature 405 (2000) 421-424; Rami, H.K., and Smith, S.A., Exp. Opin. Ther. Patents 10 (2000) 623-634 and references cited
15 therein). Three subtypes of PPARs (PPAR alpha, PPAR gamma and PPAR delta) have been identified and cloned from mouse and human. PPAR gamma, existing in three isoforms (termed PPAR gamma 1, PPAR gamma 2 and PPAR gamma 3), is the most extensively studied and is considered to be of clinical importance. The antidiabetic activity of different natural and synthetic ligands is correlated with the
20 activation of this receptor.

Thiazolidinediones are a class of compounds that selectively activate PPAR gamma and thus serve as oral insulin-sensitizing agents that lower the blood lipid and blood glucose levels. Exemplary thiazolidinediones are troglitazone, pioglitazone, ciglitazone, rosiglitazone, englitazone, BM 13.1258, BM 15.2054 and derivatives
25 thereof. The PPAR gamma activity of BM 13.1258 and BM 15.2054 has already been reported by Fürnsinn, C., et al. (Br. J. Pharmacol. 128 (1999) 1141-1148) which is incorporated by reference.

Apart from the recognised importance of PPAR gamma agonists in the area of metabolic diseases the discovery of PPAR gamma-dependent modulation of the cell
30 cycle has led to a substantial number of different approaches for the treatment of

proliferative diseases utilising compounds that bind to and thereby activate PPAR gamma. In addition to adipose tissue, PPAR gamma is reported to be highly expressed in several cancer cell lines including liposarcoma (Iijima, K., et al., Biochem. Biophys. Res. Commun. 247 (1998) 353-356), breast cancer (Mueller, E., et al., Molecular Cell (1998) 465-470; Elstner, E., et al., Proc. Nat. Acad. Sci. USA 95 (1998) 8806-8811), prostate cancer (Kubota, T., et al., Cancer Res. 58 (1998) 3344-3352) and colon cancer (Sarraf, P., et al., Nat. Med. 4 (1998) 1046-1052).

Additionally, troglitazone as a specific PPAR gamma agonist from the thiazolidinedione class is known to inhibit the growth of human cancer cells in vitro and in vivo which is disclosed in some patent applications (WO 98/25598, WO 00/18234, WO 00/30628) and described in a number of papers. In addition, thiazolidinediones including troglitazone have been shown to induce terminal differentiation in human liposarcoma cells (Tontonoz, P., et al., Proc. Nat. Acad. Sci. USA 94 (1997) 237-241). The differentiation of malignant cells represents an ideal concept for treating cancer as opposed to a cell death mediated mechanism.

The PPARs belong to type II steroid receptors that are functionally distinct from the classical steroid receptors and do not bind to their respective binding site to form a homodimer. PPAR gamma heterodimerizes with at least one other member of the steroid receptor family, the retinoid acid receptors, namely RXR alpha (Kliewer, S.A., et al., Nature 358 (1992) 771-774; Tontonoz, P., et al., Mol. Cell. Biol. 15 (1995) 351).


The combination of specific PPAR gamma ligands and RXR alpha ligands activates both receptors, leads to an additive stimulation of differentiation and results in a synergistic inhibition of the cancer cell growth (Tontonoz, P., et al., Proc. Nat. Acad. Sci. USA 94 (1997) 237-241; Elstner, E., et al., Proc. Nat. Acad. Sci. USA 95 (1998) 8806-8811).

Detailed description of the invention

The present invention is based on the unexpected finding that thiazolidinediones of formula I can be administered alone or in combination with other therapeutic agents to inhibit or reduce the growth of tumours in vivo. Activation of PPAR gamma in endothelial tissue using suitable natural or synthetic ligands is reported

to inhibit the formation of new blood vessels and thereby inhibiting angiogenesis. By reducing the vascularization the tumor size and growth can be inhibited or reduced.

In one aspect, the present invention relates to compounds of the general formula I:



(I)

wherein

- A is CH=CH or S;
- R is selected from naphthalenyl, thienyl or phenyl which could be mono- or disubstituted with C₁-C₃ alkyl, CF₃, C₁-C₃ alkoxy, F, Cl, Br or OH;
- R₁ is selected from H or C₁-C₆ alkyl;
- X is selected from S, O or NR'₂ where R' refers to H or C₁-C₆ alkyl;
- Y is CH or N;
- n is an integer from 1-3.

The enantiomers of compounds of the general formula I, their diastereomers, racemates and mixtures thereof are also included in the present invention as well as physiologically tolerated salts and solvates of these compounds with pharmaceutically acceptable, non-toxic inorganic and organic acids and bases.

Compounds of the general formula I have already been disclosed in other patent applications (WO 94/27995 and WO 98/42704) which are incorporated by reference.

Preferred compounds of formula I are those in which A is an ethylene group or sulfur, R is thienyl, unsubstituted or monosubstituted phenyl, R1 is methyl, W is oxygen, X is oxygen, Y is nitrogen and n is 2.

The most preferred compounds are:

- 5 (1) 5-{4-[2-(5-Methyl-2-phenyloxazol-4-yl)-ethoxy]-benzo[b]thiophen-7-ylmethyl}-thiazolidine-2,4-dione = BM 13.1258
- (2) 5-{4-[2-(5-Methyl-2-(thien-2-yl)oxazol-4-yl)-ethoxy]-benzo[b]thiophen-7-ylmethyl}-thiazolidine-2,4-dione = BM 15.2054
- (3) 5-{4-[2-(5-Methyl-2-phenyloxazol-4-yl)-ethoxy]-naphth-1-ylmethyl}-
10 thiazolidine-2,4-dione
- (4) 5-{4-[2-(2-(4-Fluorophenyl)-5-methyloxazol-4-yl)-ethoxy]-benzo[b]thiophen-7-ylmethyl}-thiazolidine-2,4-dione
- (5) 5-{4-[2-(2-(4-Chlorophenyl)-5-methyloxazol-4-yl)-ethoxy]-benzo[b]thiophen-7-ylmethyl}-thiazolidine-2,4-dione
- 15 (6) 5-{4-[2-(5-Methyl-2-(4-trifluoromethylphenyl)-oxazol-4-yl)-ethoxy]-benzo[b]thiophen-7-ylmethyl}-thiazolidine-2,4-dione
- (7) 5-{4-[2-(2-(4-Hydroxyphenyl)-5-methyloxazol-4-yl)-ethoxy]-benzo[b]thiophen-7-ylmethyl}-thiazolidine-2,4-dione
- (8) 5-{4-[2-(5-Methyl-2-(thien-2-yl)-oxazol-4-yl)-ethoxy]-naphthalen-1-
20 ylmethyl}-thiazolidine-2,4-dione.

Examples (7) and (8) which are novel and not disclosed in WO 94/27995 or WO 98/42704 have been prepared according to standard synthesis strategies described within these patent applications.

25 Example 7: m.p. = 127-132 °C (decomp.)

Example 8: m.p. = 158-159 °C

In another aspect, the present invention relates to the administration of a thiazolidinedione of formula I and an RXR ligand in a synergistic manner.

30 A wide variety of natural and synthetic RXR ligands as for example all-trans-retinoic acid, 9-cis-retinoic acid, phytanic acid, fenretinide, tazarotene and other derivatives of retinoic acid are known compounds that are described and disclosed

in various papers and patents and thus appropriate to be used as RXR agonists in the method of the present invention.

5 In order to produce pharmaceutical acceptable dosage forms, the compound of formula I and optionally an RXR agonist are mixed in a known manner with suitable pharmaceutical carrier substances, aromatics, flavouring and dyes and are formed for example into tablets or coated tablets or they are suspended or dissolved in water or an oil such as e.g. olive oil in addition to appropriate auxiliary substances.

10 The compound of formula I and optionally the RXR agonist can be administered orally or parenterally in a liquid or solid form. Water is preferably used as the medium that contains the stabilizing agents, solubilizers and/or buffers which are usually used for injection solutions. Such additives are for example tartrate or borate buffers, ethanol, dimethylsulfoxide, complexing agents (such as ethylenediaminetetraacetic acid), high molecular polymers (such as liquid
15 polyethylene oxide) for regulation of the viscosity or polyethylene derivatives of sorbitol anhydrides.

Solid carrier substances are e.g. starch, lactose, mannitol, methylcellulose, talcum, highly dispersed silicic acid, higher molecular fatty acids (such as stearic acid), gelatine, agar-agar, calcium phosphate, magnesium stearate, animal and vegetable
20 fats or solid high polymers (such as polyethylene glycols). Suitable formulations for the oral route can if desired contain flavourings and sweeteners.

The compound of formula I and optionally the RXR agonist are suitable to be administered also intravenously or by intramuscular, intraperitoneal, subcutaneous, intra-articular, intrasynovial, intrathecal, topical, intratumoral,
25 peritumoral, intralesional, perilesional or inhalation routes.

The administered dose for the prevention or treatment of a disease depends on the age, the health and the weight of the patient, the extent of the disease, the type of treatments which are possibly being carried out concurrently, the frequency of treatment and the type of the desired effect. The daily dose of the active compound
30 is usually 0.1 to 200 mg/kg body weight. Normally 0.5 to 100 mg/kg/day and preferably 1 to 50 mg/kg/day in one or several applications per day are effective in

order to obtain the desired results. The compound of formula I and the RXR agonist are administered as a mixture of therapeutic agents with a typical ratio of 100:1 to 1:100 and a preferable ratio of 10:1 to 1:10.

5 The compounds of formula I and optionally the RXR agonist are useful in treatment of various diseases and disorders that have been discovered to be a result of neoplastic cell proliferation. Such states include, for example, breast cancer, colon cancer, prostate cancer and liposarcoma.

10 In yet another aspect, the method of the present invention involves the use of compounds of the general formula I in combination with one or more other anti-tumor agents such as: cisplatin, carboplatin, cyclophosphamide, docetaxel, paclitaxel, bleomycin, 5-fluorouracil, 5'-deoxy-5-fluoro-N-pentyloxycarbonyl-cytidine, doxorubicine or tamoxifen.

15 The main purpose for using a compound of formula I and a conventional antineoplastic compound is to lower the doses of the potentially toxic chemotherapeutic agent while still causing tumor regression. Thiazolidinediones in general and compounds of formula I are known to be non-toxic substances. Thus, the application of a compound of formula I in addition to a conventional antineoplastic compound is expected to require up to a 100 fold decreased dosage of the toxic chemotherapeutic agent for similar efficacy compared to the
20 chemotherapeutic agent alone.

Pharmacological test results

25 It has been found that compounds of formula I, exemplary for BM 13.1258 and BM 15.2054, inhibit cell proliferation in human cancer cells of various tissue origins, e.g. in breast cancer, colon cancer and prostate cancer. Furthermore, it was demonstrated that the compounds of formula I are able to halt tumour growth in human xenograft models. Surprisingly, it was found that BM 13.1258 and BM 15.2054 reduce the metastatic load in these models, whereas troglitazone did not show this effect.

30 Female SCID beige mice were inoculated subcutaneously with 1×10^6 CX-1 human colon carcinoma cells. For 42 days 10 animals per group were orally treated with

troglitazone, BM 13.1258 or BM 15.2054 (suspension in 0.5% methylcellulose). The development of tumor volume (measured in mm³) was determined on day 45. Furthermore, following autopsy the lungs were histologically inspected for micrometastases. Inhibition of tumor growth and inhibition of number of
 5 pulmonal metastases were calculated as 100 minus tumor volume or number of metastases of the test group divided by that of the vehicle group expressed as a percentage. PPAR gamma stimulating activity of the compounds alone and in combination with the RXR specific agonist 9-cis retinoic acid were determined in accordance with a known method (Fürnsinn, C., et al., Br. J. Pharmacol. 128 (1999)
 10 1141-1148). The relative activation of PPAR-mediated gene expression is given as x-fold stimulation compared to DMSO-treated cells (activation = 1).

As demonstrated by Table 1 administration of the thiazolidinediones BM 13.1258 is effective in reducing the size of colon carcinoma in SCID beige mice. This anti-tumor effect is superior to that of troglitazone.

Table 1:

Average tumor volume and inhibition compared to vehicle treated controls

Treatment groups	Dose [mg/kg]	Mean tumor volume [mm ³]	Inhibition relative to vehicle group
Vehicle	-	1003	-
BM 15.2054	10	702	30%
BM 15.2054	50	719	28%
BM 13.1258	50	604	40%
Troglitazone	200	804	20%

Table 2 shows that by BM 15.2054 the number of lung metastases is significantly (p<0.05) reduced compared to the vehicle group. Troglitazone treatment on the
 20 contrary did not diminish the number of lung metastases.

- 8 -

Table 2:
Number of metastases per group and inhibition
compared to vehicle treated controls

Treatment groups	Dose [mg/kg]	Number of metastases	Inhibition relative to vehicle group
Vehicle	-	326	-
BM 15.2054	10	216	34%
BM 15.2054	50	223	32%
Troglitazone	200	352	-8%

5 Table 3 demonstrates that the combination of compounds of formula I with the RXR agonist 9-cis retinoic acid results in a 4-to 7-fold higher stimulation of PPAR gamma-mediated gene expression than BM 13.1258 or BM 15.2054 alone. This observed effect is not only additive, but highly synergistic.

Table 3:
Synergistic activation of PPAR gamma
in combination with 9-cis retinoic acid (RA)

	no thiazolidinedione	BM 13.1258		BM 15.2054	
		10 ⁻⁷ M	10 ⁻⁶ M	10 ⁻⁷ M	10 ⁻⁶ M
no 9-cis RA	1	3.7	4.0	4.3	3.5
9-cis RA (10 ⁻⁶ M)	4.1	24.4	28.0	15.6	18.5


10

List of References

- Elstner, E., et al., Proc. Nat. Acad. Sci. USA 95 (1998) 8806-8811
- Fürnsinn, C., et al., Br. J. Pharmacol. 128 (1999) 1141-1148
- Iijima, K., et al., Biochem. Biophys. Res. Commun. 247 (1998) 353-356
- 5 Kersten, S., et al., Nature 405 (2000) 421-424
- Kliwer, S.A., et al., Nature 358 (1992) 771-774
- Kubota, T., et al., Cancer Res. 58 (1998) 3344-3352
- Mueller, E., et al., Molecular Cell (1998) 465-470
- Pershad Singh, H.A., Exp. Opin. Invest. Drugs 8 (1999) 1859-1872
- 10 Rami, H.K., and Smith, S.A., Exp. Opin. Ther. Patents 10 (2000) 623-634
- Sarraf, P., et al., Nat. Med. 4 (1998) 1046-1052
- Tontonoz, P., et al., Mol. Cell. Biol. 15 (1995) 351
- Tontonoz, P., et al., Proc. Nat. Acad. Sci. USA 94 (1997) 237-241
- Willson, T.M., et al., J. Med. Chem. 43 (2000) 527-550
- 15 WO 00/18234
- WO 00/30628
- WO 94/27995
- WO 98/25598
- WO 98/42704
- 20

Patent Claims

1. Use of compounds of formula I



(I)

5 wherein

- A is CH=CH or S;
- R is selected from naphthalenyl, thienyl or phenyl which could be mono- or disubstituted with C₁-C₃ alkyl, CF₃, C₁-C₃ alkoxy, F, Cl, Br or OH;
- 10 - R1 is selected from H or C₁-C₆ alkyl;
- X is selected from S, O or NR'₂ where R' refers to H or C₁-C₆ alkyl;
- Y is CH or N;
- n is an integer from 1-3,

15 the enantiomers thereof, the diastereomers, racemates and mixtures as well as salts of these compounds with pharmaceutically acceptable acids and bases,

for the preparation of medicaments for inhibiting or reducing the growth of tumours.

20

2. Use of a compound according to claim 1 selected from the group consisting of

5- $\{4-[2-(5\text{-Methyl-2-phenyloxazol-4-yl})\text{-ethoxy}]\text{-benzo}[b]\text{thiophen-7-ylmethyl}\}$ -thiazolidine-2,4-dione;

5- $\{4-[2-(5\text{-Methyl-2-(thien-2-yl)oxazol-4-yl})\text{-ethoxy}]\text{-benzo}[b]\text{thiophen-7-ylmethyl}\}$ -thiazolidine-2,4-dione;

5- $\{4-[2-(5\text{-Methyl-2-phenyloxazol-4-yl})\text{-ethoxy}]\text{-naphth-1-ylmethyl}\}$ -thiazolidine-2,4-dione;

5- $\{4-[2-(2-(4\text{-Fluorophenyl})\text{-5-methyloxazol-4-yl})\text{-ethoxy}]\text{-benzo}[b]\text{thiophen-7-ylmethyl}\}$ -thiazolidine-2,4-dione;

5- $\{4-[2-(2-(4\text{-Chlorophenyl})\text{-5-methyloxazol-4-yl})\text{-ethoxy}]\text{-benzo}[b]\text{thiophen-7-ylmethyl}\}$ -thiazolidine-2,4-dione;

5- $\{4-[2-(5\text{-Methyl-2-(4-trifluoromethylphenyl)oxazol-4-yl})\text{-ethoxy}]\text{-benzo}[b]\text{thiophen-7-ylmethyl}\}$ -thiazolidine-2,4-dione;

5- $\{4-[2-(2-(4\text{-Hydroxyphenyl})\text{-5-methyloxazol-4-yl})\text{-ethoxy}]\text{-benzo}[b]\text{thiophen-7-ylmethyl}\}$ -thiazolidine-2,4-dione;

5- $\{4-[2-(5\text{-Methyl-2-(thien-2-yl)oxazol-4-yl})\text{-ethoxy}]\text{-naphthalen-1-ylmethyl}\}$ -thiazolidine-2,4-dione.

3. Use of a compound according to claim 1 or 2 in combination with an RXR agonist.

4. Use of compound according to claim 3 whereas the RXR agonist is 9-cis-retinoic acid.

5. Use of compound according to claim 1 or 2 in combination with an anti-tumour agent.

6. Use of a compound according to claim 5 wherein the antitumour agent is selected from the group consisting of cisplatin, carboplatin, cyclophosphamide, docetaxel, paclitaxel, bleomycin, 5-fluorouracil, 5'-deoxy-5-fluoro-N-pentyloxycarbonyl-cytidine, doxorubicine or tamoxifen.

7. Use of a compound of formula I according to claim 1 for the preparation of medicaments for inhibiting or reducing metastases.

8. Pharmaceutical compositions containing a compound of formula I according to claim 1 and an RXR agonist in admixture with common carriers and pharmaceutical excipients wherein the active ingredients are present in a ratio of 100:1 to 1:100.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 02/03746

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K31/425 A61P35/00 C07D417/12 C07D417/14

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, BIOSIS, FSTA

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 00 30628 A (GENENTECH INC) 2 June 2000 (2000-06-02) cited in the application abstract page 6, line 38 page 22, line 24 -page 23, line 6 claims	1-8
Y	WO 99 48529 A (AVIRAM MICHAEL ;WARNER LAMBERT CO (US); GONG BANG QIANG (US); ZHU) 30 September 1999 (1999-09-30) abstract page 3, line 7,8 claims	1-8
	--- -/--	

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *B* document member of the same patent family

Date of the actual completion of the international search	Date of mailing of the international search report
12 August 2002	30/08/2002
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016	Authorized officer Skjöldebrand, C

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 02/03746

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 98 25598 A (DANA FARBER CANCER INST INC ;ALTIOK SONER (US); SERRAF PASHA (US);) 18 June 1998 (1998-06-18) cited in the application abstract page 4, line 18-27 page 21, line 12 -page 22, line 7 page 22, line 13,14 page 23, line 3-13 claims	1-8
Y	----- FUERNSINN C ET AL: "CHRONIC AND ACUTE EFFECTS OF THIAZOLIDINEDIONES BM13.1258 AND BM15.2054 ON RAT SKELETAL MUSCLE GLUCOSE METABOLISM" BRITISH JOURNAL OF PHARMACOLOGY, BASINGSTOKE, HANTS, GB, vol. 128, no. 6, 1999, pages 1141-1148, XP001017891 ISSN: 0007-1188 cited in the application abstract	1-8
A	----- WO 94 27995 A (BOEHRINGER MANNHEIM GMBH ;MERTENS ALFRED (DE); WOLFF HANS PETER (D) 8 December 1994 (1994-12-08) cited in the application the whole document	1-8
A	----- WO 98 42704 A (WITTE ERNST CHRISTIAN ;BOEHRINGER MANNHEIM GMBH (DE); KUEHNLE HANS) 1 October 1998 (1998-10-01) cited in the application the whole document	1-8
A	----- WO 01 79202 A (HOFFMANN LA ROCHE) 25 October 2001 (2001-10-25) the whole document -----	1-8

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 02/03746

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0030628	A	02-06-2000	AU 1740900 A EP 1143953 A2 WO 0030628 A2 US 2001036955 A1	13-06-2000 17-10-2001 02-06-2000 01-11-2001
WO 9948529	A	30-09-1999	AU 1707599 A WO 9948529 A1	18-10-1999 30-09-1999
WO 9825598	A	18-06-1998	US 2002006950 A1 AU 5601898 A EP 0948324 A2 JP 2001510462 T WO 9825598 A2	17-01-2002 03-07-1998 13-10-1999 31-07-2001 18-06-1998
WO 9427995	A	08-12-1994	DE 4317320 A1 AU 682699 B2 AU 6970494 A CA 2163028 A1 CN 1124488 A ,B CZ 9502999 A3 WO 9427995 A1 EP 0700397 A1 FI 955685 A HU 75099 A2 JP 8510456 T JP 3162721 B2 KR 243783 B1 NO 954762 A NZ 267410 A PL 311752 A1 RO 114328 B1 RU 2122002 C1 SK 146195 A3 US 5599826 A	01-12-1994 16-10-1997 20-12-1994 08-12-1994 12-06-1996 13-03-1996 08-12-1994 13-03-1996 24-11-1995 28-04-1997 05-11-1996 08-05-2001 02-03-2000 24-11-1995 20-12-1996 18-03-1996 30-03-1999 20-11-1998 05-06-1996 04-02-1997
WO 9842704	A	01-10-1998	DE 19711616 A1 AU 726048 B2 AU 7036198 A BR 9808029 A CN 1250448 T WO 9842704 A1 EP 0970077 A1 HU 0001551 A2 JP 2000512312 T NO 994505 A NZ 337608 A PL 335903 A1 RU 2181724 C2 TR 9902197 T2 US 6258832 B1 US 2001008902 A1 ZA 9802326 A	24-09-1998 26-10-2000 20-10-1998 08-03-2000 12-04-2000 01-10-1998 12-01-2000 28-11-2000 19-09-2000 17-11-1999 25-05-2001 22-05-2000 27-04-2002 21-12-1999 10-07-2001 19-07-2001 20-09-1999
WO 0179202	A	25-10-2001	AU 5830601 A WO 0179202 A1 US 2001049445 A1	30-10-2001 25-10-2001 06-12-2001